

TABLE B-19

2012 VERMONT RESIDENT BIRTHS

MOTHER'S SMOKING STATUS BY COUNTY OF RESIDENCE
NUMBER OF BIRTHS AND ROW PERCENTS

| COUNTY OF RESIDENCE | NUMBER OF BIRTHS | | | TOTAL |
|---------------------|----------------------------|------------------------|---------|-------|
| | NON-SMOKERS ⁽¹⁾ | SMOKERS ⁽²⁾ | UNKNOWN | |
| ADDISON | 274 | 44 | 1 | 319 |
| BENNINGTON | 224 | 110 | 2 | 336 |
| CALEDONIA | 221 | 57 | 0 | 278 |
| CHITTENDEN | 1404 | 177 | 4 | 1585 |
| ESSEX | 40 | 8 | 0 | 48 |
| FRANKLIN | 469 | 111 | 1 | 581 |
| GRAND ISLE | 44 | 14 | 1 | 59 |
| LAMOILLE | 209 | 45 | 1 | 255 |
| ORANGE | 197 | 53 | 2 | 252 |
| ORLEANS | 214 | 56 | 0 | 270 |
| RUTLAND | 418 | 131 | 6 | 555 |
| WASHINGTON | 500 | 69 | 1 | 570 |
| WINDHAM | 288 | 94 | 23 | 405 |
| WINDSOR | 403 | 84 | 7 | 494 |
| STATE TOTAL | 4905 | 1053 | 49 | 6007 |

ROW PERCENTS - EXCLUDING UNKNOWNNS

| COUNTY OF RESIDENCE | NON-SMOKERS ⁽¹⁾ | SMOKERS ⁽²⁾ | QUITTERS ⁽³⁾ |
|---------------------|----------------------------|------------------------|-------------------------|
| ADDISON | 86.2 | 13.8 | 35.2 |
| BENNINGTON | 67.1 | 32.9 | 34.2 |
| CALEDONIA | 79.5 | 20.5 | 21.5 |
| CHITTENDEN | 88.8 | 11.2 | 22.6 |
| ESSEX | 83.3 | 16.7 | 33.3 |
| FRANKLIN | 80.9 | 19.1 | 33.3 |
| GRAND ISLE | 75.9 | 24.1 | 25.0 |
| LAMOILLE | 82.3 | 17.7 | 12.5 |
| ORANGE | 78.8 | 21.2 | 20.6 |
| ORLEANS | 79.3 | 20.7 | 35.6 |
| RUTLAND | 76.1 | 23.9 | 29.9 |
| WASHINGTON | 87.9 | 12.1 | 21.3 |
| WINDHAM | 75.4 | 24.6 | 31.0 |
| WINDSOR | 82.8 | 17.2 | 33.6 |
| STATE TOTAL | 82.3 | 17.7 | 28.4 |

SMOKING AS A RISK FACTOR FOR LOW BIRTH WEIGHT
PERCENT OF LOW BIRTH WEIGHT BABIES BY MOTHER'S SMOKING STATUS

| | NON-SMOKERS ⁽¹⁾ | SMOKERS ⁽²⁾ | QUITTERS ⁽³⁾ | ALL MOTHERS |
|----------------------|----------------------------|------------------------|-------------------------|-------------|
| PERCENT LBW (<2500g) | 5.1 | 11.5 | 5.3 | 6.2 |

⁽¹⁾ MOTHERS WHO DID NOT SMOKE CIGARETTES DURING PREGNANCY.⁽²⁾ MOTHERS WHO SMOKED CIGARETTES DURING PREGNANCY.⁽³⁾ MOTHERS WHO SMOKED CIGARETTES DURING THE THREE MONTHS BEFORE PREGNANCY OR DURING THE 1ST TRIMESTER OF PREGNANCY, BUT DID NOT SMOKE DURING THE 2ND OR 3RD TRIMESTERS OF PREGNANCY.